

Eculizumab (Soliris)

Provider Order Form



PATIENT INFORMATION

Date:	Patient Name:	DOB:
ICD-10 code (required):	ICD-10 description:	
<input type="checkbox"/> NKDA Allergies:	Weight lbs/kg:	
Patient Status: <input type="checkbox"/> New to Therapy <input type="checkbox"/> Continuing Therapy	Next Due Date (if applicable):	

PROVIDER INFORMATION

Ordering Provider:	Provider NPI:		
Referring Practice Name:	Phone:	Fax:	
Practice Address:	City:	State:	Zip Code:

NURSING

- Provide nursing care per IVX Standard Nursing Procedures, including reaction management and post-procedure observation
- Meningococcal vaccination (both conjugate and serogroup B) are required prior to initiating Soliris infusions.
- Check here if patient has already received vaccines. Fax or attach documentation of administered vaccines.
- Check here for IVX to administer vaccines as outlined below.

MENINGITIS VACCINE - Patients are required to receive first dose of both the Conjugate and serogroup b vaccines prior to initiating soliris infusions.

Unless otherwise noted, vaccines will be given 2 weeks prior to starting Soliris infusion. IVX will schedule the patient for vaccine visit followed by Soliris two weeks later. If **urgent** Soliris therapy is indicated in an unvaccinated patient, IVX Health will administer meningococcal vaccine(s) as soon as possible including same day as Soliris infusion. Additionally, provider **must prescribe** patients with 2 weeks of antibacterial drug prophylaxis.

- Check here if this is an **urgent** start.

IVX WILL ADMINISTER BOTH VACCINES AS OUTLINED BELOW.

Meningococcal conjugate (MenACWY) vaccine

(Patient will be given either Menactra or Menveo vaccine based on availability and will receive **two doses separate by at least eight weeks**. Menactra and Menveo are not interchangeable and patient will receive same product for all doses in a series.)

Serogroup B Meningococcal (MenB) vaccine

(Patient will be given Bexsero or Trumenba vaccine based on availability and will receive either the two-dose series Bexsero at least one month apart or three-dose series Trumenba at 0, 1-2, and 6 months. Bexsero and Trumenba are not interchangeable and patient will receive same product for all doses in a series.)

LABORATORY ORDERS

- CBC at each dose every _____
- CMP at each dose every _____
- Other: _____

PRE-MEDICATION ORDERS

- acetaminophen (Tylenol) 500mg / 650mg / 1000mg PO
- cetirizine (Zyrtec) 10mg PO
- loratadine (Claritin) 10mg PO
- diphenhydramine (Benadryl) 25mg / 50mg PO / IV
- methylprednisolone (Solu-Medrol) 40mg / 125mg IV
- Other: _____
- Dose: _____ Route: _____
- Frequency: _____

THERAPY ADMINISTRATION

- Eculizumab** (Soliris) in 0.9% sodium chloride, intravenous infusion
 - Dose: Induction: (Choose one. If patient has already completed induction dose, proceed to maintenance dose.)
 - 600mg weekly for the first four weeks followed by 900mg for the fifth dose one week later, then 900mg two weeks later
 - 900mg weekly for the first four weeks followed by 1200mg for the fifth dose one week later, then 1200mg two weeks later
 - Dose: Maintenance: (Choose one)
 - 900mg every two weeks 1200mg every two weeks
 - Dilute with 0.9% NS to a final concentration of 5mg/ml. (300mg doses final volume 60ml, 600mg doses final volume 120ml, 900mg doses final volume 180ml, 1200mg doses final volume 240ml.)
 - Infuse over 35 minutes in adults and 1-4 hours in pediatric patients
- Flush with 0.9% sodium chloride at the completion of infusion
- Patient is required to stay for 60-minute observation post infusion/injection
- Patient is NOT required to stay for observation time
- Refills: Zero / for 12 months / _____ (if not indicated order will expire one year from date signed)

Monitor the patient for at least one hour following completion of the infusion for signs or symptoms of an infusion reaction.

Ordering Provider: Initial here _____ and proceed to the next page.

ADULT REACTION MANAGEMENT PROTOCOL

- Observe for **hypersensitivity reaction**: Fever, chills, rigors, pruritus, rash, cough, sneezing, throat irritation, nausea, vomiting.
- If reaction occurs:
 - If indicated, stop infusion.
 - Maintain/establish vascular access.
 - IVX Health clinicians have the following PRN medications available for the following reactions.
 - Headache, pain, fever >100.4F, chills or rigors- Acetaminophen 650mg PO or Ibuprofen 400mg PO.
 - Rhinitis, allergies, hives, pruritis and other nonspecific symptoms of allergic reaction - Loratadine 10mg PO or Diphenhydramine 25-50mg PO or IV
 - Nausea, vomiting, heartburn, acid reflux- Ondansetron 4mg ODT (may repeat x 1 in 20 minutes if nausea continues, max dose 8mg) or Famotidine 20mg PO.
 - Severe Nausea, vomiting, heartburn, acid reflux- Ondansetron 4mg SIVP (may repeat x 1 in 20 minutes if nausea continues, max dose 8mg) or Famotidine 20mg SIVP.
 - Hypotension (90/60), vasovagal response- Place patient in reclined position, administer 0.9% Sodium Chloride IV 500ml. May repeat to keep BP >90/60, maximum of 1000ml, monitor vital signs.
 - Hypertension (>30 mmHg increase from baseline or >180 mmHg SBP): Clonidine 0.1mg and wait 45 minutes, may administer Amlodipine 5mg if hypertension persists
 - Chest pain/discomfort, shortness of breath- Oxygen 2-15 liters, titrate to keep Spo2 >92%.
 - Famotidine 20mg IV- Refractory to other treatments given
 - Solumedrol 125mg IV- Refractory to other treatments given.
 - When symptoms resolve resume infusion at 50% previous rate and increase per manufactures guidelines.
 - Notify referring provider as clinically appropriate and follow clinical escalation protocol.
- Severe allergic/anaphylactic reaction:**
 - If symptoms are rapidly progressing or continuing after administration of prn medications above and signs symptoms of severe allergic/anaphylactic reaction (angioedema, swelling of the mouth, tongue, lips, or airway, dyspnea, bronchospasm with or without hypotension or hypertension).
 - Call 911.
 - Initiate basic life support as needed.
 - Bring the **AED** to the patient (Attach pads if indicated).
 - **Epinephrine**- administer 0.3mg of a 1:1,000 (1mg/ml) concentration intramuscularly (preferably outer thigh), may be repeated every 5-15 minutes as needed to a maximum of 3 doses.
 - Place patient in recumbent position, elevate lower extremities.
 - **Oxygen**- administer 2-15 liters/minute or 100 percent oxygen as needed maintain SpO2 >92 percent.
 - **IV Fluids**- Treat hypotension with normal saline bolus of 500ml, repeat as needed to maintain systolic BP >90.
 - Administer **diphenhydramine** 50mg IV or Famotidine 20mg IVP, if not previously given.
 - Administer **methylprednisolone** 125mg IVP, if not previously given.
 - Continuous monitoring of blood pressure, pulse oximetry, and heart rate.
 - Notify clinical executive, DON or CMO, when appropriate. Must be done same day. Do not delay treatment.

Patient Name	Patient Date of Birth
Provider Name (Print)	
Provider Signature	Date

Email ivxintake@ivxhealth.com or fax this form, insurance card (both sides), demographics, recent H&P, labs, and supporting clinicals to:

TAMPA: 844-946-0849 ___Brandon ___Carrollwood ___Wesley Chapel ___St. Pete's PHILADELPHIA: 844-820-9641 ___Malvern ___Bensalem ___Montgomeryville BAY AREA: 844-889-0275 ___San Mateo ___Fremont ___San Ramon ___Sunnyvale CHICAGO: 312-253-7244 ___Glenview ___Schaumburg ___Lombard ___Naperville KANSAS CITY: 844-900-1292 ___Overland Park ___Lee's Summit ___Briarcliff INDIANAPOLIS: 844-983-2028 ___Emerson Pointe ___Fishers Corner ___Park Meridian	ORLANDO: 844-946-0867 ___Altamonte Springs ___Waterford Lakes HARRISBURG: 844-859-4235 ___East Shore ___West Shore COLUMBUS: 844-627-2675 ___Dublin ___Pickerington ___Grove City CINCINNATI: 844-946-0868 ___Colerain ___Hyde Park ___Union Centre NASHVILLE: 844-627-2518 ___Brentwood ___Hendersonville
---	---

Precision is now a part of IVX Health. To refer to a Precision center, fax to 888-615-1445: ___Donelson ___Cool Springs
 ___Clarksville ___Murfreesboro ___Knoxville ___Chattanooga ___Morristown ___Collierville ___Jackson ___Memphis ___Lowell, AR