

Idursulfase (Elaprase)

Provider Order Form



PATIENT INFORMATION

Date:	Patient Name:	DOB:
ICD-10 code (required):	ICD-10 description:	
<input type="checkbox"/> NKDA Allergies:	Weight lbs/kg:	
Patient Status: <input type="checkbox"/> New to Therapy <input type="checkbox"/> Continuing Therapy	Next Due Date (if applicable):	

PROVIDER INFORMATION

Ordering Provider:	Provider NPI:		
Referring Practice Name:	Phone:	Fax:	
Practice Address:	City:	State:	Zip Code:

NURSING

- Provide nursing care per IVX Standard Nursing Procedures, including reaction management

LABORATORY ORDERS

- CBC at each dose every _____
- CMP at each dose every _____
- CRP at each dose every _____
- Other: _____

PRE-MEDICATION ORDERS

- acetaminophen (Tylenol) 500mg / 650mg / 1000mg PO
- cetirizine (Zyrtec) 10mg PO
- loratadine (Claritin) 10mg PO
- diphenhydramine (Benadryl) 25mg / 50mg PO / IV
- methylprednisolone (Solu-Medrol) 40mg / 125mg IV
- Other: _____
- Dose: _____ Route: _____
- Frequency: _____

THERAPY ADMINISTRATION

- Idursulfase** (Elaprase) in 100ml 0.9% sodium chloride, intravenous infusion
 - Dose: 0.5mg/kg
 - Route: intravenous
 - Frequency: once every week
 - The total volume of infusion should be administered over a period of 3 hours, which may be gradually reduced to 1 hour if no hypersensitivity reactions are observed.
- Infuse with a low-protein-binding 0.2 micrometer (μ m) in-line filter.
- Flush with 0.9% sodium chloride at the completion of infusion
- Patient is required to stay for 30-minute observation post infusion
- Patient is NOT required to stay for observation time
- Refills: Zero / for 12 months / _____ (if not indicated order will expire one year from date signed)

SPECIAL INSTRUCTIONS

Ordering Provider: Initial here _____ and proceed to the next page.

ADULT REACTION MANAGEMENT PROTOCOL

- Observe for **hypersensitivity reaction**: Fever, chills, rigors, pruritus, rash, cough, sneezing, throat irritation, nausea, vomiting.
- If reaction occurs:
 - If indicated, stop infusion.
 - Maintain/establish vascular access.
 - IVX Health clinicians have the following PRN medications available for the following reactions.
 - Headache, pain, fever >100.4F, chills or rigors- Acetaminophen 650mg PO or Ibuprofen 400mg PO.
 - Rhinitis, allergies, hives, pruritis and other nonspecific symptoms of allergic reaction - Loratadine 10mg PO or Diphenhydramine 25-50mg PO or IV
 - Nausea, vomiting, heartburn, acid reflux- Ondansetron 4mg ODT (may repeat x 1 in 20 minutes if nausea continues, max dose 8mg) or Famotidine 20mg PO.
 - Severe Nausea, vomiting, heartburn, acid reflux- Ondansetron 4mg SIVP (may repeat x 1 in 20 minutes if nausea continues, max dose 8mg) or Famotidine 20mg SIVP.
 - Hypotension (90/60), vasovagal response- Place patient in reclined position, administer 0.9% Sodium Chloride IV 500ml. May repeat to keep BP >90/60, maximum of 1000ml, monitor vital signs.
 - Hypertension (>30 mmHg increase from baseline or >180 mmHg SBP): Clonidine 0.1mg and wait 45 minutes, may administer Amlodipine 5mg if hypertension persists
 - Chest pain/discomfort, shortness of breath- Oxygen 2-15 liters, titrate to keep Spo2 >92%.
 - Famotidine 20mg IV- Refractory to other treatments given
 - Solumedrol 125mg IV- Refractory to other treatments given.
 - When symptoms resolve resume infusion at 50% previous rate and increase per manufactures guidelines.
 - Notify referring provider as clinically appropriate and follow clinical escalation protocol.
- Severe allergic/anaphylactic reaction:**
 - If symptoms are rapidly progressing or continuing after administration of prn medications above and signs symptoms of severe allergic/anaphylactic reaction (angioedema, swelling of the mouth, tongue, lips, or airway, dyspnea, bronchospasm with or without hypotension or hypertension).
 - Call 911.
 - Initiate basic life support as needed.
 - Bring the **AED** to the patient (Attach pads if indicated).
 - **Epinephrine**- administer 0.3mg of a 1:1,000 (1mg/ml) concentration intramuscularly (preferably outer thigh), may be repeated every 5-15 minutes as needed to a maximum of 3 doses.
 - Place patient in recumbent position, elevate lower extremities.
 - **Oxygen**- administer 2-15 liters/minute or 100 percent oxygen as needed maintain SpO2 >92 percent.
 - **IV Fluids**- Treat hypotension with normal saline bolus of 500ml, repeat as needed to maintain systolic BP >90.
 - Administer **diphenhydramine** 50mg IV or Famotidine 20mg IVP, if not previously given.
 - Administer **methylprednisolone** 125mg IVP, if not previously given.
 - Continuous monitoring of blood pressure, pulse oximetry, and heart rate.
 - Notify clinical executive, DON or CMO, when appropriate. Must be done same day. Do not delay treatment.

Patient Name

Patient Date of Birth

Provider Name (Print)

Provider Signature

Date

Email ivxintake@ivxhealth.com or fax this form, insurance card (both sides), demographics, recent H&P, labs, and supporting clinicals to:

TAMPA: 844-946-0849 ___Brandon ___Carrollwood ___Wesley Chapel ___St. Pete's

ORLANDO: 844-946-0867 ___Altamonte Springs ___Waterford Lakes

PHILADELPHIA: 844-820-9641 ___Malvern ___Bensalem ___Montgomeryville

HARRISBURG: 844-859-4235 ___East Shore ___West Shore

BAY AREA: 844-889-0275 ___San Mateo ___Fremont ___San Ramon ___Sunnyvale

COLUMBUS: 844-627-2675 ___Dublin ___Pickerington ___Grove City

CHICAGO: 312-253-7244 ___Glenview ___Schaumburg ___Lombard ___Naperville

CINCINNATI: 844-946-0868 ___Colerain ___Hyde Park ___Union Centre

KANSAS CITY: 844-900-1292 ___Overland Park ___Lee's Summit ___Briarcliff

NASHVILLE: 844-627-2518 ___Brentwood ___Hendersonville

INDIANAPOLIS: 844-983-2028 ___Emerson Pointe ___Fishers Corner ___Park Meridian



Precision is now a part of IVX Health. To refer to a Precision center, fax to 888-615-1445: ___Donelson ___Cool Springs

___Clarksville ___Murfreesboro ___Knoxville ___Chattanooga ___Morristown ___Collierville ___Jackson ___Memphis ___Lowell, AR